

Investigating the impact of remediation efforts on Hediste diversicolor in the Seine estuary using multiple levels of biological organization

Andrew Barrick, Iris Barjhoux, Jean-Marie Marion, Amélie Châtel, Hanane Perrein-Ettajani, Mohammed Mouloud, Patrick Gillet, Isabelle Métais,

Catherine Mouneyrac

▶ To cite this version:

Andrew Barrick, Iris Barjhoux, Jean-Marie Marion, Amélie Châtel, Hanane Perrein-Ettajani, et al.. Investigating the impact of remediation efforts on Hediste diversicolor in the Seine estuary using multiple levels of biological organization. Marine Environmental Research, 2022, 173, pp.105528. 10.1016/j.marenvres.2021.105528. hal-03689804

HAL Id: hal-03689804 https://uco.hal.science/hal-03689804

Submitted on 5 Jan 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

Investigating the Impact of Remediation Efforts on Hediste diversicolor in the Seine Estuary Using Multiple Levels of Biological Organization Stuary Using Multiple Levels of Biological Organization

A.Barrick^{a,b}, I. Barjhoux^c, J.M. Marion^d, A. Châtel^a, H. Perrien-Ettajani^a, M. Mouloud^a, P. Gillet^a, I.
 Mettais^a, C. Mouneyrac^a

- a Mer Molécules Sante (MMS), Université Catholique de l'Ouest, 3 Place André Leroy, BP10808,
 49008 Angers Cedex 01, France
- 8 b Cawthron Institute, 98 Halifax st East, 7010, Nelson, New Zealand

9 c UMR-I 02 Stress Environnementaux et Biosurveillance des milieu aquatiques (SEBIO), UFR SEN,

10 Université de Reims Champagne-Ardenne, Moulin de la Housse, BP 1039, Reims, France

11

12 d LARIS (Laboratoire Angevin de Recherche en Ingénierie des Systèmes), EA-7315, Université

- 13 Catholique de l'Ouest, 3 Place André Leroy, BP10808, 49008 Angers Cedex 01, France
- 14

15 Abstract

16 The Seine Estuary is historically one of the most contaminated estuaries in Europe. In 2002 the 17 estuary underwent major redevelopment which emphasized environmental remediation to measure 18 biological effects. The present study investigated the health status of the sentinel species Hediste 19 diversicolor in the Seine estuary by comparing data from a sampling campaign immediately after the 20 Seine underwent reconstruction and 10 years afterward. Both studies implemented multiple levels of 21 biological organization ranging from enzymatic biomarkers to population density. Integrative 22 modeling was used to establish a holistic status assessment for H. diversicolor in the Seine. H. 23 diversicolor populations in the Seine estuary were impacted in both campaigns suggesting no 24 significant improvements over the 10-year period. This interpretation however can be improved by 25 increasing the number of reference sites used to establish the baseline natural variation, integrating 26 additional environmental parameters into the model and measuring additional biological endpoints, 27 such as reproductive status.

- 28
- 29
- 29
- 30
- 31
- 32
- 33
- 55
- 34
- 35
- 36
- 37
- 38

39 Introduction

40 The Seine Basin in France represents a catchment area of approximately 78,650 km² and 41 encompasses 40% of the French economic activity including intensive agricultural activities, strong 42 industrialization, and significant urbanization (Billen et al., 2007). Due to a long history of human 43 activity, the Seine estuary is considered one of the most contaminated water systems in Europe with 44 a wide variety of persistent pollutants in the environment including metals, polycyclic aromatic 45 hydrocarbons (PAHs), and polychlorinated biphenyls (PCBs) (Blanchard et al., 2007; Hamzeh et al., 46 2014; Thévenot et al., 2007). This results in a complex mixture of anthropogenic contamination that 47 arises in short-term and long-term impacts on ecosystem health and makes effective mitigation 48 difficult. The Seine estuary is a major water system that is crucial for the life history of many 49 invertebrates, fish, and birds, making it crucial to investigate the severity of anthropogenic impacts 50 making anthropogenic contaminants an ecological concern. Starting in 2002, the estuary underwent 51 major redevelopment and rehabilitation work (dredging to improve water circulation, reconstruction 52 of meanders, prolongation of breakwaters, and creation of an artificial island) under the Port 2000 53 project which extended the harbor and closed out the northern flank of the estuary (Tecchio et al., 54 2016). The construction of the port fundamentally shifted hydrology and deposition of sediments 55 into the ecosystem which may have mixed effects on organisms (Dauvin et al., 2006). Sediments are 56 a sink for numerous persistent compounds and reworking sediments could potentially reintroduce 57 sequestered contaminants into the ecosystem and threaten both benthic and pelagic organisms 58 (Fisson, 2014). The project to expand the port integrated ecological considerations in an attempt to 59 mitigate anthropogenic impacts but major changes in hydrology and sedimentary characteristics of 60 the estuary caused clear modifications in benthic communities (Cuvilliez et al., 2009; Dauvin et al., 61 2010).

62 Investigating environmental impact of such a development plan, it is functionally and 63 economically unrealistic to screen for every potential contaminant, and implementing biological tools 64 may be more cost-effective (Amiard and Amiard-Triquet, 2008). To assess the health status of the 65 Seine estuary it was proposed to use biomarkers addressing multiple levels of biological organization 66 from sub-individual endpoints to population dynamics, as biomarkers may provide early warning 67 tools to identify stress caused by contaminants (Amiard and Amiard-Triquet, 2008). The effectiveness 68 of biomarkers in biomonitoring frameworks is recognized but their deployment is limited risk 69 assessment strategies due the wide range of responses that can be observed and the effect this has 70 on interpretation of impact. Concentrations of anthropogenic contaminants can be determined by 71 chemical analysis to identify if harmful compounds are present at concentrations above the natural 72 background making it comparatively easy, when coupled with biological response, to identify 73 whether or not a site is contaminated and if amelioration efforts improve environmental conditions.

74 To assess the influence of the Port 2000 construction on the Seine estuary, the National 75 Ecotoxicology Program (PNETOX) conducted an environmental risk assessment survey during the 76 construction of the harbour (2002-2004). The project implemented a multitude of biotic and abiotic 77 endpoints and compared values to the Authie estuary. At the time of the initial study, Authie was 78 identified to be an estuary with similar environmental parameters, equivalent latitude and 79 comparatively lower levels of contamination (Durou et al., 2008). Industrialization is also absent in 80 the Authie estuary but low levels of metal and PAH contaminant are present and, in the context of 81 the study, was considered an ideal comparator for assessing the quality Seine estuary (Francescangeli 82 et al., 2020). Hediste diversicolor (H. diversicolor), a sediment-dwelling polychaete worm, was 83 selected for the study due to its status as a keystone, sentinel species with a broad geographic 84 distribution (Durou et al., 2007a; Kalman et al., 2009; Thit et al., 2015). The sampling survey 85 investigated the impact at different levels of biological organization on H. diversicolor (Amiard-86 Triquet and Rainbow, 2017). The results of the project indicate that biomass and density of the 87 suprabenthos declined between 2002 and 2004 (approximately 80%) and that population levels of H. 88 diversicolor diminished too (Gillet et al., 2008; Tecchio et al., 2016). The study also highlighted 89 evidence linking anthropogenic activities in the Seine estuary to sub-individual effects in H. 90 diversicolor (Durou et al., 2008; Durou and Mouneyrac, 2007). Previous studies have demonstrated 91 that benthic communities show a steep decline after major alterations and that longer periods post-92 construction are necessary to verify the effectiveness of remediation efforts (Meng et al., 2018). It 93 can also be unclear how to interpret data from multiple levels of biological organization in a clear and 94 concise interpretation of environmental health. When considering the response of organisms 95 however, it can be unclear how to interpret data from multiple levels of biological organization and 96 create clear conclusions on the health status of organisms in a polluted site.

97 To follow up on the longevity of adverse effects of pollution in the Seine estuary, project 98 ECOTONES (2015-2017) was initiated approximately 10 years later measuring the same biological 99 endpoints as PNETOX. Project ECOTONES aimed to assess if the overall health of H. diversicolor in the 100 Seine estuary improved because of remediation efforts. Interpreting organismal responses using 101 multiple levels of biological organization can often be unclear, leading to inconsistent interpretations 102 of health status in contaminated estuaries. Applying a method of biomarker response integration 103 that weights biological endpoints based on the level of biological organisation can potentially be used 104 to establish concise conclusions on health status.

105 One of the proposed methods to integrate biological responses at different levels of 106 organization is based on the use of a line of evidence (LOE) for biomarkers (Barjhoux et al., 2018; Piva 107 et al., 2011). This LOE model compiles biological responses at multiple levels of organization into an 108 integrated hazard quotient (HQ) associated with a hazard class ranking from absent (no observable 109 biological impact) to severe (significant important impact), allowing for the comparison of temporal 110 biomarker datasets through quantitative (HQ values) and qualitative (hazard classes) methods. Six 111 biological endpoints were used in the present study: oxidative stress biomarkers (catalase (Cat), 112 glutathione-s-transferase (GST), and thiobarbituric reactive substances (TBARs)), neurotoxicity-113 specific response (acetylcholinesterase activity (AChE)), energy reserves (total lipid and glycogen 114 content) and population density. For each integrated endpoint, reference level and thresholds were 115 derived from the baseline assessment criteria (BAC) and the environmental assessment criteria (EAC) 116 previously defined (Barrick et al., 2018a, 2016) which integrate the effects of confounding factors 117 (salinity, weight of individuals and temperature) on the studied biomarkers into the analysis.

118

119 Materials and methods

120 Sample site and collection

The Seine estuary in northern France is a system influenced by high tides and the estuary is split in three spatial areas: the Northern Flank (NF), the Central Channel (CC), and the Southern Flank (SF) (Figure 1). Both PNETOX and ECOTONES sampling campaigns were carried out in the NF region of the estuary. In both projects, *H. diversicolor* individuals were collected in the intertidal zone at low tide in the Seine estuary (49°27'17.52" N and 0°11'19.57" E). Historical data for the spring, summer, and fall seasons of the 2002-2004 period in the Seine estuary were collected from previous studies (Durou et al., 2007a; Durou and Mouneyrac, 2007; Gillet et al., 2008). 128 Sampling for ECOTONES was carried out two times each year (spring and summer for 2015-2017). 129 The temperature and the salinity of water remaining on the mudflat surface were recorded using a 130 field thermometer (CG867) and a salinometer (WTW LF 196). Biological endpoints, time points and 131 sampling protocols comparable to the historical data generated in PNETOX. To analyze density, four 132 quadrats (625 cm² and 25 cm deep) were collected during each sampling campaign (Ait Alla et al., 133 2006; Gillet et al., 2008). Quadrats were then sieved to collect organisms (200µm mesh size) and 134 organisms retained were collected and stored in 70% ethanol. For biomarker analyses, samples were 135 collected gently by hand in parallel with the collection of samples for density. Twenty sexually 136 immature worms of equivalent size (0.11±0.03g wet weight) were collected and individually analyzed 137 for enzymatic biomarkers (AChE, GST, Cat, and TBARs) and twenty worms of equivalent size were 138 collected and individually analyzed for energetic reserves (glycogen and lipids). Samples were 139 immediately frozen in liquid nitrogen, transported to the laboratory, and stored at -80°C until 140 analysis.

141

142 Biomarker analysis

143 Biomarkers in the present study were analyzed following protocols used in the PNETOX 144 program. Briefly, enzymatic biomarker responses were assessed using the techniques described in 145 Durou et al., 2007. Samples were individually homogenized in buffer solution (0.08 M Tris, 2 M NaCl, 146 1 mM of 1,4-Dithiothreitol, and cocktail protease inhibitor diluted to 1:1000 at pH 7.4 and 4 °C) in a 147 1:3 ratio of tissue to the solution using a mortar and pestle. The homogenates were then centrifuged 148 for 30 minutes at 9000×g. Total protein content was calculated by following the Bradford assays 149 (Bradford, 1976; Galgani and Bocquene, 1991). GST activity was determined spectroscopically at 150 340nm by measuring the rate that 1-choloro-dinitrobenzene is conjugated with reduced glutathione 151 (Habig et al., 1974). Cat activity was calculated by measuring the consumption of H_2O_2 152 spectroscopically at 240 nm. Results were expressed as µmoles.min⁻¹.mg of protein⁻¹ (Aebi, 1984). 153 AChE activity was calculated by using a coupled reaction measuring the catalyzation of 154 acetylthiocholine to produce thiocholine (which is then reduced by 5,5'-Dithiobis(2-nitrobenzoic acid) 155 to produce thionitrobenzoate (Bocquené et al., 1990). The rate of the reaction was measured by colorimetric spectroscopy. Lipid peroxidation was measured by the formation of thiobarbituric 156 157 reactive substances quantified by malondialdehyde absorbance at 530 nm (Knight et al., 1988).

Energetic reserves were analyzed following the protocol previously described in Durou et al. (2007). Individuals were ground into a powder using liquid nitrogen and then prepared as a homogenate in a citrate buffer (0.1 M, pH 5.0). Total lipid content was determined following the method defined in (Frings et al., 1972). Glycogen content was determined by partitioning the sample in two (with or without enzymatic digestion by amyloglucosidase) following the protocol described in (Carr and Neff, 1984a).

164

165 Data integration within Biomarker index and development of LOE model

To establish a baseline that accounted for natural variation, the method described in Barrick et al. (2016) was applied. Multi-polynomial regression models were previously developed for each biomarker to integrate confounding factors (salinity, temperature, and weight of individual) into the analysis, establishing thresholds of expected natural variation using data from the Authie estuary as a reference point. Authie was used as the reference point for this study to maintain comparison with the historical data and, while the use of several reference sites is ideal, there is limited comparable 172 data available for other estuaries. For each regression model, a baseline assessment criterion (BAC) 173 was used to define the expected range of variation that would occur naturally in healthy individuals. 174 A banding approach was used to define the threshold of disturbance, also called environmental 175 assessment criterion (EAC), with twice the lower limit of the BAC used for stress-induced inhibition 176 endpoints (AChE, energy reserve assays, and density) to indicate the biomarker is severely impacted. 177 Formulas were previously defined for biomarkers (AChE, GST, CAT, TBARS, glycogen, and lipids) in 178 Barrick et al. (2016, 2018). For population density, a regression model was calculated following the 179 same method. BAC/EAC values were established for each sampling periods in the Seine estuary using 180 confounding factors measured during sampling periods for both projects.

181 To characterize in situ biological responses for H. diversicolor for integrated biomarker 182 responses an LOE model was generated following a previously defined method with some 183 adjustments to fit data collected (Barjhoux et al., 2018, Piva et al., 2011). The application of the LOE 184 model requires the establishment of reference values (or control values), thresholds (Th) values 185 indicating the range of expected natural variation, and an effect threshold above which physiological 186 disturbances are expected to occur. Mean values from the BAC models were used as reference 187 values and the Th value was calculated from the lower or upper limits of the BAC depending on how 188 the biological responses change under stress conditions, i.e. the 'ecologically relevant' direction of 189 biomarker modulation. Effect threshold values were derived from EAC values for each biological 190 endpoint. In the LOE model, only unilateral modulations were taken into consideration. Details of the 191 LOE parametrization and calculations are presented in the supplemental material and summarized 192 below.

193 To further develop the model, measured responses were expressed as a percentage change 194 over the respective reference value and divided by the Th value. A correction factor was applied to 195 the resulting ratio using the following method: a 0.2-factor for values less than or equal to the Th 196 values, a 0.5-factor for responses above the Th value but lower than the effect threshold, and a 1-197 factor for values that exceeded the effect threshold. These calculations resulted in an 'effect' value 198 E(i) for each biological response which was reported on an evaluation grid for hazard classification ranging from absent to severe. E(i) values were then weighted (Ew(i)) according to the biological 199 200 significance of the considered biomarker following the approach described in Piva et al. (2011): a 201 weight of 0.75 was applied to TBARS and AChE, a 0.5-weight was assigned to CAT and GST, a 1.2 202 weight was given to lipid and glycogen, and a weight of 2.8 for density measurements as an intermediate value between mortality (3.0), histopathology and reproductive failure (2.4). 203

204 Finally, a cumulative biomarker hazard quotient (BiomHQ) was generated by calculating the 205 weighted average of the Ew(i) values of biomarkers attributed to the moderate class of hazard and 206 summing the Ew(i) values with responses belonging to the major or the severe class of hazard. The 207 contribution of each endpoint and class of endpoint to the BiomHQ value was also calculated as 208 described by Barjhoux et al. (2018). Another global index (BiomHQ index) was also generated by 209 adding the percentage of biomarkers divided into the different classes of hazard weighed with 210 coefficients increasing with the attributed hazard (from 0.7 for absent to 8 for severe; according to 211 Piva et al. 2011). This new index was reported on an evaluation grid for a global hazard classification 212 (absent to severe).

213

214 <u>Results</u>

215 When assessing the health status of *H. diversicolor* using individual biomarker hazard 216 rankings many of the biological endpoints demonstrated wide variations ranging from "absent" to "severe" between sampling campaigns (Table 1). Of the endpoints used in the present study TBARS, glycogen and density were the endpoints the indicated stress. Annually, hazard rankings for the PNETOX campaign suggested stress response for TBARS, AChE, GST, catalase, lipids, glycogen and density for 2002 whereas in 2003 AChE, GST, glycogen and density indicated stress and in 2004 only GST and density demonstrated a stress response. For project ECOTONES TBARS, AChE, lipids, glycogen and density consistently suggested a stress response for 2015-2017 with catalase indicating moderate stress in 2015.

224 Individual biological endpoints were then weighted following the approach used in Piva et al. 225 (2011) and results were integrated into biomarker hazard quotients (BiomHQ) to classify stress 226 response (Figure 2). This index represents the cumulative response profiles of all biological endpoints 227 reletative to reference values and can be interpreted as a quantitative indicator of the intensity of 228 deviation from expected values (ie out of the range of basal variations defined by the reference and 229 the threshold Th values). Results from the PNETOX program showed limited variations of the 230 BiomHQ values, with maximal values (13-15.5) observed between July 2002 and February 2003 and 231 the lowest (0-3) between November 2003 and May 2004. On the contrary, BiomHQ values calculated 232 with ECOTONES results proved to be generally higher and more variable with indices reaching about 233 20 and 35 in March 2015 and July 2015, respectively. Individual BiomHQ values are calculated based 234 on the number of biomarker responses/ severity of response for each data set. In some cases this 235 leads to lower values indicating higher stress response (e.g. 2002 vs 2003). The BiomHQ index, a 236 more qualitative description of the biological disturbance observed at each sampling time, was 237 calculated according to the weighted distribution of biomarkers within the different classes of risk 238 attributed for each E(i) value. This second index is used to attribute the risk classification for each 239 sampling period and the annual averages (see figure S1 in the supplemental material).

240 The contribution of each biomarker to the interpretation of hazard were also assessed 241 (Figure 3). In many of the sampling campaigns for PNETOX, oxidative stress (mainly TBARS and GST), 242 neurotoxicity (AChE), and populational (density) markers were major contributing factors to the 243 BiomHQ values. Other responses such as energetic reserves and CAT also influenced the level of 244 hazard in some of the PNETOX sampling campaigns but more punctually. For the ECOTONES sampling 245 campaign, TBARs were a major contributor in defining the health status for H. diversicolor the Seine 246 estuary. Energetic reserves contributed more to the hazard classification in the ECOTONES sampling 247 campaign compared to the PNETOX samples. AChE, CAT, and population density also factored into 248 the level of environmental hazard in ECOTONES but their contribution was dependent on the sample 249 period and no clear profile could be identified.

Annually, the population density was a major indicator for 2003 and 2004 whereas TBARS, neurotoxicity, and energetic reserves (in particular glycogen) were the most contributive biological endpoints for 2002 and 2015-2017. As observed for the first PNETOX campaigns, the highest BiomHQ values are associated with more complex contribution profiles indicating that impact can be detected at different levels of response.

255

256 Discussion

Polychaete worms are ideal target organisms for monitoring ecosystem health as they are sensitive to anthropogenic activity and play a crucial role as contaminant vectors across the trophic chain (Green Etxabe et al., 2021). The objective of the present study was to investigate if remediation efforts in the Seine estuary had improved the health status *of H.* diversicolor in a way that accounts for natural variatio using a multi-biomarker approach following an integrative method of analysis. 262 The BAC/EAC model attempts to account for effects environmental parameters (temperature, salinity 263 and weight) on the biological endpoints used in the study by defining the range of values expected to 264 naturally occur for each endpoint. The results of the model demonstrated large fluctuations 265 seasonally and annually for all biological endpoints, suggesting that natural variation may not be 266 completely accounted for and that longer time series may be necessary to characterize this 267 relationship more appropriately. Additional research is necessary to further clarify the complex 268 relationship between biological endpoints, natural variation and effects of anthropogenic activity 269 (Galloway, 2006). Measurement of temperature and salinity in the ECOTONES project were 270 measured to conform to the how data was collected in the PNETOX campaign and measured 271 environmental temperature and salinity at the time of sampling. Using an average temperature and 272 salinity across several timepoints might improve the model's ability to predict how these parameters 273 influence measured biological endpoints. Integrating additional parameters such as granulometry, 274 available organic matter and reproductive status can also significantly improve the model as these 275 endpoints can alter metabolic activity which, in turn, can influence subindividual endpoints (Díaz-276 Jaramillo et al., 2011).

277 Annually PNETOX data indicated less stress in 2003 and 2004 when compared to the initial 278 sampling in 2002 which would be consistent with recovery after a major environmental perturbation 279 like the construction of the port. Data for the ECOTONES campaign was relatively consistent 2015-280 2017 which suggested the site was majorly impacted. This observation would indicate that despite 281 the 10-year difference between the two projects, H. diversicolor populations in the Seine estuary 282 were still majorly impacted and effects of contamination likely persistent and chronic (please see 283 supplemental materials for data regarding measured contaminants present. The site used to collect 284 H. diversicolor in the Seine estuary is an active shipping zone and has been demonstrated to have 285 high concentrations of PAHs, PCBs and organocholorine pesticides which have been previously, and 286 proportionally, accumulated in H. diversicolor (Durou et al., 2007a; Lesueur et al., 2015). The Seine 287 estuary also has high concentrations of metal in the sediment with previous studies indicating a poor 288 correlation between metal concentrations in the environment and concentrations in H. diversicolor 289 (Amiard-Triquet et al., 2007; Berthet et al., 2015; Saiz-Salinas and Francés-Zubillaga, 1997). 290 Nonetheless, chronic exposure of *H. diversicolor* to both persistent organic pollutants and metallic 291 trace elements present in the sediment of the Seine estuary may be responsible for the effects 292 highlighted by the LOE model.

293 Biological endpoints used in the study account for a wide range of responses including 294 neurotoxicity, oxidative stress, energetic reserves, and population density. Surprisingly, endpoints 295 that consistently suggested adverse effects were mainly TBARS, glycogen and density suggesting that 296 H. diversicolor is impacted at multiple levels of biological organization. Results also indicated that, of 297 the subcellular endpoints, TBARS was one of the more consistent biomarkers. One potential 298 explanation for this observation is that TBARS are indicative of damage and suggest that defense 299 mechanisms are consistently overwhelmed in the Seine estuary, implying chronic effects. Enzymatic 300 biomarkers (AChE, GST and Cat) displayed seasonal patterns which is unsurprising as the previous BAC/EAC models indicated a wide range of natural variation (Barrick et al., 2018b; Beliaeff and 301 302 Bocquené, 2004; Hagger et al., 2006). It has also been demonstrated that enzymatic biomarkers can 303 be influenced seasonal variation in solar exposure and adjustments to energetic budgets (Díaz-304 Jaramillo et al., 2011). Future adjustments may need to consider a piece-wise integrative model in 305 order to investigate the relationship between levels of biological organization at specific time points, 306 in particular the relationship between subcellular responses and dynamic energy budgets .

307 When considering the results of energetic reserves, the LOE model suggested glycogen levels 308 are impacted in the Seine estuary, which is consistent with the BAC model previously generated 309 (Barrick et al., 2016). Glycogen also had a considerable contribution to the interpretation of 310 environmental health status suggesting that it may be a useful endpoint in identifying sites adversely 311 impacted by anthropogenic activity. With respect to lipids there was some evidence suggesting 312 moderate stress but overall was less impacted than glycogen. This may suggest that glycogen is more 313 responsive to environmental condition which may make it an ideal endpoint for measuring 314 anthropogenic activity. One potential explanation for this is that glycogen is the primary energy 315 resource (maintenance, reproduction, growth) when accounting for environmental stress (Carr and 316 Neff, 1984b). It is also important to consider that energetic reserves are highly dependent on the 317 reproductive status of the organisms and that contamination may alter biological clocks (Pook et al., 318 2009). Previous work on the PNETOX data demonstrated that the sexual reproductive status of H. 319 diversicolor in the Seine were equivalent to individuals in Authie, suggesting an allocation of energy 320 towards reproduction rather than growth (Durou et al., 2007b). Integration of reproductive status 321 into the model would be beneficial as it would help characterize the effects of this relationship on 322 other biological endpoints.

323 Population density values consistently fell outside the range of expected natural variation, 324 providing evidence that the Seine is still majorly impacted and potentially influencing the ecosystem. 325 This is consistent with a previous interpretation which suggested that low population densities may 326 be due to irrigation and sediment reworks in the Seine estuary (Gillet et al., 2008). It is however 327 important to note that polychaete population densities can display significant variations across time 328 and space (Kristensen, 1984). Despite Authie being considered a good comparator estuary for the 329 Seine intersite differences, such as nuances in sediment texture and algal cover, can potentially make 330 it an unreliable comparator site for density assessments (Alzeny et al., 2021). Integrating multiple 331 references sites and accounting for these environmental parameters can potential improve the 332 quality of the model. Perturbations to the density of H. diversicolor indicated impact during the 333 PNETOX sampling campaign may be attributed to immediate alterations to the sediment dynamics as 334 result of the construction process rather than the presence of contamination. The results of the 335 model suggested 'moderate' to 'major' effects towards H. diversicolor density across the 10-year 336 period, suggesting the site is still impacted. Previous research on macrofaunal composition in the 337 Seine estuary are resilient to anthropogenic perturbation and one potential explanation for the 338 interpretation of the model in the present study is that the response profile is localized (Bacouillard 339 et al., 2020). Future alterations to the model should consider including multiple sampling sites in the 340 Seine estuary.

341 No true reference values exist but the model attempts to account for expected environmental 342 variation through the application of a BAC/EAC index. While this approach allows for clear, concise 343 results for communicating environmental health to regulatory bodies, it is important to note that 344 additional research is necessary to clarify the complex relationship between biological endpoints, 345 natural environmental variation and effects of anthropogenic activity (Galloway, 2006). Some of the 346 endpoints used in the model demonstrated clear seasonal patterns indicating that longer time series 347 may be necessary. It also necessary to look at additional endpoints linked to seasonal variation such 348 as nutrient load, turbidity, and sediment transport.

The LOE model establishes a first step in developing a comprehensive model to provide an holistic assessment of *H. diversicolor* to improve the assessment of health status. Improvements can be made to the initial model by including other biological endpoints such as genotoxicity, reproductive success, or scope for growth. This was not done in the present study due to a lack of necessary data to develop relevant BAC indices for these endpoints. In addition to this, different biological endpoints have varying degrees of sensitivity to environmental disturbance and it may be necessary to clarify these relationships in future studies. Through the integration of population density into the LOE model it becomes possible to extrapolate how sublethal endpoints can lead to adverse effects on the population. Future research will need to expand on the relationship between endpoints as well as how environmental parameters may influence these interpretations.

359

360 <u>Conclusion</u>

"The proposed model provides a holistic assessment of the status of *H. diversicolor* individuals in 361 362 the Seine estuary but to be effective there needs to be effective integration with other LOEs, in 363 particular sediment type and concentrations of contaminants, to effectively establish a weight of 364 evidence approach that characterizes environmental quality. The present study proposes a first 365 attempt at developing an LOE model for H. diversicolor that integrates subcellular endpoints, subindividual and populational effects to establish a holistic approach to environmental health. This 366 approach was applied to biological responses analyzed in the sentinel species H. diversicolor 367 368 following two long term studies in the Seine estuary. The LOE model implemented the use of a 369 BAC/EAC model which attempts to correct for variation due to natural variations such as 370 temperature, salinity and weight of the organisms. The quality of this assessment can be improved 371 upon by integrating time series data for the environmental parameters, which could potentially 372 capture how climate change influences *H. diveriscolor*.

There are however major limitations with the results as the data demonstrates large seasonal variations in the interpretation of health status for the biomarkers and future research will need to investigate how to account for this. The use of only one reference site to establish the BAC/EAC index also limits the efficacy of the model and endeavors should be made establish additional data at other low impact sites. Finally, the interpretation of environmental impact is also hampered by the lack of of contaminant data and the results would be strengthened by chemistry data.

Overall, the present study demonstrates that H. diversicolor could be an effective model 379 380 organisms for environmental hazard assessment. The model could however be improved by 381 additional endpoints, such as genotoxicity and reproductive stage, can further describe the health of 382 individuals in the Seine estuary. Results suggested that H. diversicolor in the Seine estuary are 383 moderately to majorly impacted by environmental condition, indicating that the status of the Seine 384 estuary has not notably improved between the two sampling campaigns. This however is only one 385 line of evidence and other lines of evidence need to generate in order to effectively characterize the 386 impact human activity has had on the Seine estuary, in particular coupling additional 387 physicochemical parameters such as contamination levels in the environment and bioaccumulation 388 measurements in key species into a weight of evidence approach."

389

390 Acknowledgements

This research was funded through the public interest group Seine-Aval (GIP Seine-Aval, SA5 2014-ECONTONES) under the project ECOTONES. Samples were collected for the ECOTONES data using the generation of the model were collected with assistance from the Normandy Littoral Monitoring Unit.

394

395 **References**

- Aebi, H., 1984. Oxygen Radicals in Biological Systems. Methods Enzymol. 105, 121–126.
 https://doi.org/10.1016/S0076-6879(84)05016-3
- Ait Alla, A., Gillet, P., Deutsch, B., Moukrim, A., Bergayou, H., 2006. Response of Nereis diversicolor (
 Polychaeta , Nereidae) populations to reduced wastewater discharge in the polluted estuary of
 Oued Souss , Bay of Agadir , Morocco 70. https://doi.org/10.1016/j.ecss.2006.01.028
- Alzeny, A., El-tohamy, W.S., Branch, A., 2021. Spatio-temporal population and reproductive
 responses in Hediste diversicolor (Polychaeta: Nereididae) from the Damietta Harbor Barge
 Canal, Egypt. 25, 721–735.
- Amiard-Triquet, C., Durou, C., Amiard, J.C., Berthet, B., Billon, G., Budzinski, H., Créach, A., Roméo,
 M., Debenay, J.-P., Deloffre, J., Denis, F., Denis, L., Ferrero, T., Gillet, P., Hummel, H.,
 Mouneyrac, C., Ouddane, B., Petit, F., Poirier, L., Quillet, L., Smith, B., Sylvestre, F., 2007. A
 comprehensive methodology for the ass essment of the health status of estuarine ecosystems.
 Ices 1–17.
- Amiard-Triquet, C., Rainbow, P., 2017. Environmental Assessment of Estuarine Ecosystems: A Case
 Study, 1st ed. CRC Press: Taylor & Francis Group.
- Amiard, J.C., Amiard-Triquet, C., 2008. Les biomarqueurs dans l'évaluation de l'état écologique des
 milieux aquatiques.
- Bacouillard, L., Dauvin, J., Desroy, N., Geiger, K.J., Gentil, F., 2020. Long-term spatio-temporal
 changes of the muddy fine sand benthic community of the Bay of Seine (eastern English
 Channel) 161. https://doi.org/10.1016/j.marenvres.2020.105062
- 416 Barjhoux, I., Fechner, L., Lebrun, J., Anzil, A., Ayrault, S., Budzinski, H., Cachot, J., Charron, L.,
- 417 Chaumot, A., Clérandeau, C., Dedourge-Geffard, O., Faburé, J., Fançois, A., Geffard, O., George,
- 418 I., Labadie, P., Lévi, Y., Munoz, G., Noury, P., Oziol, L., Quéau, H., Servais, P., Uher, E., Urien, N.,
- 419 Geffard, A., 2018. Application of a multidisciplinary and integrative weight-of-evidence
- 420 approach to a 1-year monitoring survey of the Seine River. Environ. Sci. Pollut. Res. 25, 23404–
 421 23429. https://doi.org/10.1007/s11356-016-6993-6
- Barrick, A., Chatel, A., Marion, J.M., Perrein-Ettajani, H., Bruneau, M., Mouneyrac, C., 2016. A novel
 methodology for the determination of biomarker baseline levels in the marine polychaete
 Hediste diversicolor. Mar. Pollut. Bull. 108, 275–280.
- 425 https://doi.org/10.1016/j.marpolbul.2016.04.056
- Barrick, A., Marion, J., Perrein-ettajani, H., Châtel, A., Mouneyrac, C., 2018a. Baseline levels of
 biochemical biomarkers in the endobenthic ragworm Hediste diversicolor as useful tools in
 biological monitoring of estuaries under anthropogenic pressure. Mar. Pollut. Bull. 129, 81–85.
 https://doi.org/10.1016/j.marpolbul.2018.02.006
- Barrick, A., Marion, J.M., Perrein-Ettajani, H., Châtel, A., Mouneyrac, C., 2018b. Baseline levels of
 biochemical biomarkers in the endobenthic ragworm Hediste diversicolor as useful tools in
 biological monitoring of estuaries under anthropogenic pressure. Mar. Pollut. Bull. 129, 81–85.
 https://doi.org/10.1016/j.marpolbul.2018.02.006
- Beliaeff, B., Bocquené, G., 2004. Exploratory data analysis of the Mediterranean component of the
 BEEP programme. Mar. Environ. Res. 58, 239–244.
- 436 https://doi.org/10.1016/j.marenvres.2004.03.065
- Beliaeff, B., Burgeot, T., 2002. Integrated biomarker response: A useful tool for ecological risk
 assessment. Environ. Toxicol. Chem. 21, 1316–1322. https://doi.org/10.1002/etc.5620210629

- Berthet, B., Mouneyrac, C., Amiard, J.C., Berthelot, Y., Hen, A. Le, Mastain, O., 2015. Accumulation
 and Soluble Binding of Cadmium , Copper , and Zinc in the Polychaete Hediste diversicolor from
 Coastal Sites with Different Trace Metal E nvironmental C ontamination Accumulation and
 Soluble Binding of Cadmium , Copper , and Zinc in the Polyc. https://doi.org/10.1007/s00244-
- 443 003-0135-0
- Billen, G., Garnier, J., Mouchel, J.M., Silvestre, M., 2007. The Seine system: Introduction to a
 multidisciplinary approach of the functioning of a regional river system. Sci. Total Environ. 375,
 1–12. https://doi.org/10.1016/j.scitotenv.2006.12.001
- Blanchard, M., Teil, M.J., Guigon, E., Larcher-Tiphagne, K., Ollivon, D., Garban, B., Chevreuil, M., 2007.
 Persistent toxic substance inputs to the river Seine basin (France) via atmospheric deposition
 and urban sludge application. Sci. Total Environ. 375, 232–243.
 https://doi.org/10.1016/j.scitotenv.2006.12.012
- 451 Bocquené, G., Galgani, F., Truquet, P., 1990. Characterisation and assay conditions for use of AChE 452 activity from several marine species in pollution monitoring. Mar. Environ. Res. 30, 75–89.
- Bradford, M.M., 1976. A rapid and sensitive method for the quantitation of microgram quantities of
 protein utilizing the principle of protein-dye binding. Anal. Biochem. 72, 248–254.
 https://doi.org/10.1016/0003-2697(76)90527-3
- 456 Carr, R.S., Neff, J.M., 1984a. Quantitative semi-automated enzymatic assay for tissue glycogen.
 457 Comp. Biochem. Physiol. -- Part B Biochem. 77, 447–449. https://doi.org/10.1016/0305458 0491(84)90258-X
- 459 Carr, R.S., Neff, J.M., 1984b. Field assessment of biochemical stress indices for the sandworm
 460 Neanthes virens (Sars). Mar. Environ. Res. 14, 267–279. https://doi.org/10.1016/0141461 1136(84)90082-5
- 462 Cuvilliez, A., Deloffre, J., La, R., Bessineton, C., 2009. Geomorphology Morphological responses of an
 463 estuarine intertidal mud fl at to constructions since 1978 to 2005 : The Seine estuary (France)
 464 104, 165–174. https://doi.org/10.1016/j.geomorph.2008.08.010
- Damiens, G., Gnassia-Barelli, M., Loquès, F., Roméo, M., Salbert, V., 2007. Integrated biomarker
 response index as a useful tool for environmental assessment evaluated using transplanted
 mussels. Chemosphere 66, 574–583. https://doi.org/10.1016/j.chemosphere.2006.05.032
- 468 Dauvin, J., Bellan, G., Bellan-santini, D., 2010. Benthic indicators : From subjectivity to objectivity –
 469 Where is the line ? Mar. Pollut. Bull. 60, 947–953.
 470 https://doi.org/10.1016/j.marpolbul.2010.03.028
- 471 Dauvin, J.C., Desroy, N., Janson, A.L., Vallet, C., Duhamel, S., 2006. Recent changes in estuarine
 472 benthic and suprabenthic communities resulting from the development of harbour
 473 infrastructure. Mar. Pollut. Bull. 53, 80–90. https://doi.org/10.1016/j.marpolbul.2005.09.020
- 474 Díaz-Jaramillo, M., Martins da Rocha, A., Gomes, V., Bianchini, A., Monserrat, J.M., Sáez, K., Barra, R.,
 475 2011. Multibiomarker approach at different organization levels in the estuarine Perinereis
 476 gualpensis (Polychaeta; Nereididae) under chronic and acute pollution conditions. Sci. Total
 477 Environ. 410–411, 126–135. https://doi.org/10.1016/j.scitotenv.2011.09.007
- 478 Durou, C., Mouneyrac, C., 2007. Linking steroid hormone levels to sexual maturity index and energy
 479 reserves in Nereis diversicolor from clean and polluted estuaries. Gen. Comp. Endocrinol. 150,
 480 106–113. https://doi.org/10.1016/j.ygcen.2006.07.019
- 481 Durou, C., Mouneyrac, C., Amiard-triquet, C., 2008. Environmental quality assessment in estuarine
 482 ecosystems : Use of biometric measurements and fecundity of the ragworm Nereis diversicolor

- 483 (Polychaeta , Nereididae) 42, 2157–2165. https://doi.org/10.1016/j.watres.2007.11.028
- 484 Durou, C., Poirier, L., Amiard, J.C., Budzinski, H., Gnassia-Barelli, M., Lemenach, K., Peluhet, L.,
 485 Mouneyrac, C., Roméo, M., Amiard-Triquet, C., 2007a. Biomonitoring in a clean and a multi486 contaminated estuary based on biomarkers and chemical analyses in the endobenthic worm
 487 Nereis diversicolor. Environ. Pollut. 148, 445–458.
- 488 https://doi.org/10.1016/j.envpol.2006.12.022
- 489 Durou, C., Smith, B.D., Mouneyrac, C., Mouloud, M., Gnassia-barelli, M., Gillet, P., Deutch, B., Amiard 490 triquet, C., 2007b. From biomarkers to population responses in Nereis diversicolor : Assessment
- 491 of stress in estuarine ecosystems 66, 402–411. https://doi.org/10.1016/j.ecoenv.2006.02.016
- 492 Fisson, C., 2014. Qualité des eaux de l'estuaire de la Seine.
- Francescangeli, F., Quijada, M., Armynot du Châtelet, E., Frontalini, F., Trentesaux, A., Billon, G.,
 Bouchet, V.M.P., 2020. Multidisciplinary study to monitor consequences of pollution on
 intertidal benthic ecosystems (Hauts de France, English Channel, France): Comparison with
 natural areas. Mar. Environ. Res. 160. https://doi.org/10.1016/j.marenvres.2020.105034
- 497 Frings, C.S., Fendley, T.W., Dunn, R.T., Queen, C.A., 1972. ImprovedDeterminationof Total Serum
 498 Lipids by the Sulfo-Phospho-Vanillin Reaction 18, 673–674.

Galgani, F., Bocquene, G., 1991. Semi-automated colorimetric and enzymatic assays for aquatic
organisms using microplate readers. Water Res. 25, 147–150. https://doi.org/10.1016/00431354(91)90023-J

- Galloway, T.S., 2006. Biomarkers in environmental and human health risk assessment. Mar. Pollut.
 Bull. 53, 606–613. https://doi.org/10.1016/j.marpolbul.2006.08.013
- Gillet, P., Mouloud, M., Durou, C., Deutsch, B., 2008. Response of Nereis diversicolor population (
 Polychaeta , Nereididae) to the pollution impact e Authie and Seine estuaries (France) 76,
 201–210. https://doi.org/10.1016/j.ecss.2007.07.004
- Green Etxabe, A., Pini, J.M., Short, S., Cunha, L., Kille, P., Watson, G.J., 2021. Identifying conserved
 polychaete molecular markers of metal exposure: Comparative analyses using the Alitta virens
 (Annelida, Lophotrochozoa) transcriptome. Comp. Biochem. Physiol. Part C Toxicol.
 Pharmacol. 240, 108913. https://doi.org/10.1016/j.cbpc.2020.108913
- Habig, W., Pabst, M., Jakoby, W., 1974. Glutathione S-Transferases. Joural Biol. Chem. 249, 7130–
 7139.
- Hagger, J.A., Jones, À.M.B., Leonard, À.P., Owen, R., À, T.S.G., 2006. Biomarkers and Integrated
 Environmental Risk Assessment : Are There More Questions Than Answers ? Integr. Environ.
 Assess. Manag. 2, 312–329.
- Hamzeh, M., Ouddane, B., Daye, M., 2014. Trace Metal Mobilization from Surficial Sediments of the
 Seine River Estuary Trace Metal Mobilization from Surficial Sediments of the Seine River Estuary
 0–15. https://doi.org/10.1007/s11270-014-1878-0
- Kalman, J., Palais, F., Amiard, J.C., Mouneyrac, C., Muntz, A., Blasco, J., Riba, I., Amiard-Triquet, C.,
 2009. Assessment of the health status of populations of the ragworm Nereis diversicolor using
 biomarkers at different levels of biological organisation. Mar. Ecol. Prog. Ser. 393, 55–67.
 https://doi.org/10.3354/meps08239
- Knight, J.A., Robert, K., Mcclellan, L., 1988. Specificity of the thiobarbituric acid reaction: its use in
 studies of lipid peroxidation. Clin. Chem. 34, 2433–8.
- 525 Kristensen, E., 1984. Life cycle, growth and production in estuarine populations of the polychaetes

- 526
 Nereis virens and N. diversicolor. Holarct. Ecol. 7, 249–256. https://doi.org/10.1111/j.1600

 527
 0587.1984.tb01128.x
- Lesueur, T., Boulangé-lecomte, C., Restoux, G., Deloffre, J., Xuereb, B., Le, K., Budzinski, H.,
 Petrucciani, N., Marie, S., Petit, F., Forget-leray, J., 2015. Ecotoxicology and Environmental
 Safety Toxicity of sediment-bound pollutants in the Seine estuary , France , using a Eurytemora
 af fi nis larval bioassay. Ecotoxicol. Environ. Saf. 113, 169–175.
- 532 https://doi.org/10.1016/j.ecoenv.2014.11.033
- Meng, X., Jiang, X., Li, Z., Wang, J., Cooper, K.M., Xie, Z., 2018. Responses of macroinvertebrates and
 local environment to short-term commercial sand dredging practices in a flood-plain lake. Sci.
 Total Environ. 631–632, 1350–1359. https://doi.org/10.1016/j.scitotenv.2018.03.086
- Piva, F., Ciaprini, F., Onorati, F., Benedetti, M., Fattorini, D., Ausili, A., Regoli, F., 2011. Assessing
 sediment hazard through a weight of evidence approach with bioindicator organisms: A
 practical model to elaborate data from sediment chemistry, bioavailability, biomarkers and
 ecotoxicological bioassays. Chemosphere 83, 475–485.
- 540 https://doi.org/10.1016/j.chemosphere.2010.12.064
- 541 Pook, C., Lewis, C., Galloway, T., 2009. The metabolic and fitness costs associated with metal
- resistance in Nereis diversicolor. Mar. Pollut. Bull. 58, 1063–1071.
- 543 https://doi.org/10.1016/j.marpolbul.2009.02.003
- Saiz-Salinas, J., Francés-Zubillaga, G., 1997. Nereis diversicolor : an Unreliable Biomonitor of Metal
 Contamination in the ' Ria de Bilbao ' (Spain). Mar. Ecol. 18, 113–125.

Tecchio, S., Chaalali, A., Raoux, A., Tous, A., Lequesne, J., Girardin, V., Lassalle, G., Cachera, M., Riou,
P., Lobry, J., Dauvin, J., Niquil, N., 2016. Evaluating ecosystem-level anthropogenic impacts in a
stressed transitional environment : The case of the Seine estuary. Ecol. Indic. 61, 833–845.
https://doi.org/10.1016/j.ecolind.2015.10.036

- Thévenot, D.R., Moilleron, R., Lestel, L., Gromaire, M.C., Rocher, V., Cambier, P., Bonté, P., Colin, J.L.,
 de Pontevès, C., Meybeck, M., 2007. Critical budget of metal sources and pathways in the Seine
 River basin (1994-2003) for Cd, Cr, Cu, Hg, Ni, Pb and Zn. Sci. Total Environ. 375, 180–203.
 https://doi.org/10.1016/j.scitotenv.2006.12.008
- Thit, A., Banta, G.T., Selck, H., 2015. Bioaccumulation, subcellular distribution and toxicity of
 sediment-associated copper in the ragworm Nereis diversicolor: The relative importance of
 aqueous copper, copper oxide nanoparticles and microparticles. Environ. Pollut. 202, 50–57.
 https://doi.org/10.1016/j.envpol.2015.02.025
- 558



Figure 1: Location of the study area in the Seine estuary – Northern flank. Black bar indicates 1km. (Credits: Google Maps, Images @2019 Google, Data SIO, NOAA, U.S. Navyn NGA, GEBCO, Images @2019 CNES, Airbus, Landsat, Copernicus, Maxar Technologies, Données cartographique @2019).



Figure 2 : Evaluation and classification of an integrated assessment of biological endpoints measured in the Seine estuary. Values are based on the biomarker hazard quotient (BiomHQ) and indicate the degree in which biological endpoints deviated from the range of expected natural variation.



Figure 3: Contribution of each biomarker to the BiomHQ value calculated for each sampling campaign. N/A indicates that contributions could not be calculated due to BiomHQ value equals to 0.



Figure 4: Contribution of each class of biological endpoints to the BiomHQ value calculated for each sampling campaign. N/A indicates that contributions could not be calculated due to BiomHQ value equals to 0.

Program	Year	Month	TBARS	AChE	GST	Catalase	Lipids	Glycogen	Density
PNETOX	2002	Feb.	Severe	Severe	Major	Major	Slight	Absent	Slight
		Apr.	Severe	Absent	Absent	Severe	Slight	Major	Absent
		Jul.	Severe	Severe	Absent	Absent	Absent	Slight	Major
		Sept.	Severe	Absent	Absent	Absent	Major	Major	Major
	2003	Feb.	Absent	Severe	Major	Absent	Slight	Absent	Severe
		May	Absent	Absent	Absent	Absent	Absent	Absent	Major
		Aug.	Slight	Absent	Absent	Absent	Absent	Major	Major
		Nov.	Absent	Absent	Major	Slight	Absent	Absent	Slight
	2004	Feb.	Absent	Major	Major	Absent	Slight	Absent	Absent
		May	Absent	Absent	Absent	Absent	Absent	Absent	Slight
		Aug.	Absent	Absent	Absent	Absent	Absent	Major	Major
		Nov.	Absent	Absent	Absent	Absent	Absent	Absent	Severe
ECOTONES	2015	Mar.	Severe	Absent	Absent	Major	Major	Slight	Absent
		Jun.	N/A	Major	Absent	Absent	Absent	Absent	Major
	2016	Mar.	Severe	Absent	Absent	Absent	Major	Major	Absent
		Jul.	Severe	Severe	Absent	Absent	Absent	Major	Major
	2017	Mar.	Severe	Absent	Absent	Absent	Major	Major	Absent
		Jun.	Major	Severe	Absent	Absent	Absent	Major	Major
ANNUAL AVERAGES	2002		Severe	Major	Absent	Moderate	Slight	Moderate	Moderate
	2003		Absent	Slight	Moderate	Absent	Absent	Slight	Major
	2004		Absent	Absent	Slight	Absent	Absent	Absent	Moderate
	2015		Severe	Moderate	Absent	Moderate	Moderate	Slight	Moderate
	2016		Severe	Moderate	Absent	Absent	Moderate	Major	Moderate
	2017		Severe	Moderate	Absent	Absent	Moderate	Major	Moderate

Table 1: Classification of hazard for biological endpoints within the Biomarker LOE model, based on the calculated effect (E(i)) values. Results are presented based upon each sampling campaign and annually.